

K X-ray satellite relative intensities of Ca excited by photons

M V R Murti¹, B Mallikarjuna Rao², B Seetharami Reddy² and K Parthasaradhi³

¹Centre of Studies in Resources Engineering, IIT, Mumbai-400 076

²Department of Nuclear Physics, Andhra University, Visakhapatnam-530 003, India

³Medical Physics Division, North Western University Medical School, Evanston Hospital, Evanston, U.S.A.

Received 1 June 1998, accepted 24 November 1998

Abstract The intensity ratios of $K_{\alpha}L^1/K_{\alpha}L^0$ and $K_{\beta}L^1/K_{\beta}L^0$ of Ca in CaCO_3 and $\text{CaHPO}_4 \cdot 2\text{H}_2\text{O}$ compounds are measured by photon (Cr tube) excitation employing a plane crystal (LiF 200) wavelength dispersive spectrometer. Comparison of these ratios with the published data shows a slight evidence of the dependence of the ratio on the incident photon excitation energy and chemical composition of the Ca compounds and a definite dependence of the ratios on multiple ionization.

Keywords K X-ray satellites, calcium compounds, photon excitation

PACS No. 33.20.Rm

Introduction

Single K-shell ionization produces the normal K_{α} and K_{β} characteristic X-rays from atoms. Single K plus L shell ionization produces K X-ray satellites. These are designated as KL^n where n is the number of L holes. This multiple ionization can be produced by photons, electrons, ions and heavier ions [1-6]. However, as the mode of excitation changes from photons to ions to heavier ions the probability of producing more number of L holes (multiple ionization) increases. With photon excitation, the first satellite can well be studied because of the following advantages: (1) non-deterioration of the target and (2) negligible secondary excitations. A few attempts [3, 7, 8] have been made by photon excitation particularly below $Z = 17$. Recently Murthy *et al* [9] carried out studies on the relative intensity ratios of the first satellites of K_{α} and K_{β} , respectively to those of K_{α} and K_{β} characteristic X-rays emitted with single K hole of Ca in CaCO_3 . They carried out measurements using a plane crystal (LiF 200) wavelength dispersive spectrometer with X-rays produced from Rh anode. In the present communication, we have studied the relative intensity ratios of the first satellites of K_{α} and K_{β} X-rays to those of their respective K X-rays emitted with single K hole (diagram lines) of Ca in CaCO_3 and

Correspondence to : B Mallikarjuna Rao

$\text{CaHPO}_4 \cdot 2\text{H}_2\text{O}$ compounds using Cr tube X-rays for excitation with a view to see (1) the dependence of these ratios on the excitation energy and (2) the possible evidence of chemical effects.

The measurements are carried out with a Philips 1410 wavelength dispersive crystal ($\text{LiF } 200$; $2d = 4.027 \text{ \AA}$) spectrometer with Cr tube X-rays at 30 kV voltage and 30mA current. Murthy *et al* [9] used settings of 40 kV and 30 mA for the Rh tube. In our case the tube voltage is lesser and also the energy of the characteristic X-rays emitted by Cr is lower than the energy of X-rays emitted by Rh. Pellets of 5-mm thick and 50 mm diameter of pure powders of CaCO_3 and $\text{CaHPO}_4 \cdot 2\text{H}_2\text{O}$ are used. The data are collected in 2 theta steps of 0.05° intervals. The measured spectra with the various X-ray lines are shown in the Figure 1. The intensity ratios of




Figure 1. K X-ray spectra of Ca excited by photons. The counts of CaHPO_4 are increased by an order of 10 to avoid overlap of the spectra

$K_\alpha L^1 / K_\alpha L^0$ and $K_\beta L^1 / K_\beta L^0$ are evaluated from their respective areas and are given in Table 1 along with the available data [3, 6, 9, 10]. Within the brackets, the corresponding values reported by Murthy *et al* [9] are also given in the same table. The ratio by electron excitation of Parratt [6] is in excellent agreement with that of Murthy *et al* [9]. However, the present values are higher than those of Murthy *et al* [9]. This shows some evidence for the dependence of the ratio on the excitation energy. As the excitation energy decreases the ratio is found to be increasing. Secondly, the relative intensities of the two compounds studied show slight evidence of possible chemical effects. The ion excitation values are higher than the present values.

Table 1. K X ray satellite relative intensities (%)

Ratio	Present work (Cr tube) photon excitation		Parrott [6] electron excitation	McWherter <i>et al</i> [3] proton excitation	Aberg [10] Theory
	CaCO_3	$\text{CaHPO}_4 \cdot 2\text{H}_2\text{O}$	CaO	Ca	
$K_\alpha L^1 / K_\alpha L^0 [K_{\alpha(1,2)}]$	2.8 ± 0.2 (2.2 ± 0.2)*	2.6 ± 0.2	2.2	10.0	3.1
$K_\beta L^1 / K_\beta L^0 [K_{\beta(1,3)}]$	1.1 ± 0.2 (0.8 ± 0.2)*	1.4 ± 0.2		4.8	

Values in the brackets are of Murthy *et al* [9]

*Used Rh tube anode

which shows that the ratios also depend definitely on the probability of multiple ionization, which increases for ion- and heavier ion-excitations. However, Aberg's [10] theoretical values based on Ne like structure, with no screening, are in better agreement with present measurements than with the previous values [6, 9].

Acknowledgment

This work has been performed under the DST collaborative research project between the Centre of Studies in Resources Engineering, IIT, Mumbai and the Department of Nuclear Physics, Andhra University, Visakhapatnam. The authors gratefully acknowledge the financial assistance given by the Department of Science & Technology, Government of India.

References

- [1] O Keski-Rahkonen, K Reinikainen and E Mikkola *Phys. Scr.* **28** 179 (1983)
- [2] R L Watson, B Bandong, J M Sanders and K Parthasaradhi *Phys. Scr.* **31** 184 (1985)
- [3] G Ramesh Babu, K Parthasaradhi, M L N Raju, M V R Murti and K S Rao *Physica* **138C** 201 (1986)
- [4] D K Olsen and C F More *Phys. Rev. Lett.* **33** 194 (1974)
- [5] J McWherter, J Bolger, C F More and P Richards *Z. Phys.* **263** 283 (1973)
- [6] L G Parrott *Phys. Rev.* **50** 1 (1936)
- [7] J Utrianen, M Linkoaho, E Rantavuori, T Aberg and G Graeffe *Z. Naturforsch.* **23A** 1178 (1968)
- [8] K Parthasaradhi, G Ramesh Babu, V R K Murthy, M V R Murti and K S Rao *Nucl. Instrum. Meth.* **A255** 54 (1987)
- [9] V R K Murthy, K Parthasaradhi and M V R Murti *X-Ray Spectrometry* **27** 23 (1998)
- [10] T Aberg *Phys. Lett.* **26** 515 (1968)